

Modelling and computing homotopy types: I *

Ronald Brown

December 21, 2016

Abstract

The aim of this article is to explain a philosophy for applying higher dimensional Seifert-van Kampen Theorems, and how the use of groupoids and strict higher groupoids resolves some foundational anomalies in algebraic topology at the border between homology and homotopy. We explain some applications to filtered spaces, and special cases of them, while a sequel will show the relevance to n -cubes of pointed spaces.

Contents

Introduction	2
1 Statement of the philosophy	4
2 Broad and Narrow Algebraic Data	4
3 Five Anomalies in Algebraic Topology	6
4 The origins of algebraic topology	7
5 Enter groupoids	8
6 Higher dimensions?	11

*This article has developed from presentations given at Liverpool, Durham, Kansas SU, Chicago, IHP, Galway, Aveiro, whom I thank for their hospitality. It has been submitted for publication in the special issue, forthcoming in 2017, of *Indagationes Mathematicae* honoring the memory of L. E. J. Brouwer.

7	Need also “Commutative cubes”	13
8	Cubical sets in algebraic topology	16
9	Strict homotopy double groupoids?	16
10	Crossed Modules	17
11	Groupoids in higher homotopy theory?	18
12	Filtered spaces	21
13	Crossed complexes and their classifying space	22
14	Why filtered spaces?	23

Introduction

The usual algebraic topology approach to homotopy types is to aim first to calculate specific invariants and then look for some way of amalgamating these to determine a homotopy type. A standard complication is that in a space, identifications in low dimensions affect high dimensional homotopy invariants.

The philosophy which we state in outline in Section [1](#) was developed since about 1965 and was in response to a gradual realisation of certain anomalies in traditional algebraic topology which we explain in Section [3](#).

A basic reason for formulating and trying to analyse these issues is given by Einstein in the correspondence published in [[Ein90](#)]:

... the following questions must burningly interest me as a disciple of science: What goal will be reached by the science to which I am dedicating myself? What is essential and what is based only on the accidents of development? ... Concepts which have proved useful for ordering things easily assume so great an authority over us, that we forget their terrestrial origin and accept them as unalterable facts. ... It is therefore not just an idle game to exercise our ability to analyse familiar concepts, and to demonstrate the conditions on which their justification and usefulness depend, and the way in which these developed, little by little ...

One conclusion from progress made with resolving some of these anomalies is the difficulty of

dealing with “bare” topological spaces, i.e. spaces with no additional structure. In order to calculate something about a space X we need some data on X and this data will usually have some structure. Thus our algebraic invariants need to recognise such structure. Further, in view of the influence of low dimensions on high dimensions, we need algebraic invariants which have structure in a range of dimensions. In particular, the single base point which is necessary to define the n th homotopy group is a rather limited kind of structure.

A traditional view on a prospective Seifert-van Kampen theorem in dimension 2 would be that we need to determine the Postnikov invariant $k_X \in H^3(\pi_1(X), \pi_2(X))$ when X is a union. So we need $\pi_1(X)$ by the standard Seifert-van Kampen Theorem; then find $\pi_2(X)$ of a union; then find k_X in terms of the invariants of the pieces. The last two steps seem impossible.

However if X is a CW-complex, then by the main result of [MLW50], k_X is determined by the crossed module

$$\pi_2(X, X^1, x) \rightarrow \pi_1(X^1, x).$$

The advantage of this approach is that crossed modules form an algebraic structure for which colimits exist and can in principle be calculated, and further they can be thought of as structures in dimensions 1 and 2. So we can at least formulate the possibility that this crossed module of a union could be the colimit of the crossed modules of the pieces. What is much more difficult is to see how to prove, or disprove, such a result.

A key to the Mac Lane-Whitehead paper was the result in [Whi49] that the crossed module

$$\pi_2(X \cup \{e_\lambda^2\}, X, x) \rightarrow \pi_1(X, x)$$

could be described as a free crossed module¹ on the characteristic maps of the 2-cells e_λ^2 , $\lambda \in \Lambda$. Thus we see a universal property in 2-dimensional relative homotopy theory. So the authors of [BH78] began in 1974 to look for proving a 2-dimensional van Kampen theorem in a *relative* situation, i.e. one with more structure; see the Strategic Analysis 9.1.

The intuition that there might be a 2-dimensional van Kampen Theorem came in 1965 with an idea for the use of forms of double groupoids. We explain more on this idea in Section 6.

A further feature of interest to me has been that the van Kampen theorem is a *local-to-global theorem*, and such theorems are of wide importance in mathematics and science. Further, the standard van Kampen theorem for groups is a nonabelian theorem of this type, which is unusual. Algebraic models which could allow a higher dimensional version have the possibility of being really new. Such a view seemed therefore well worth pursuing, although it has been termed “idiosyncratic”. It can now be seen as related to issues² discussed in [Ati02]. Any dichotomy between geometry and algebra as suggested in that paper is softened by the methods given in Section 2. Tools of category theory are also needed for the effective application of the theory.

¹This result is sometimes stated but rarely proved in texts on algebraic topology.

²local to global, increase in dimensions, commutative to noncommutative, linear to non linear, geometry versus algebra

1 Statement of the philosophy

The general philosophy which has developed is to consider categories and functors

$$\left(\begin{array}{c} \text{Topological} \\ \text{Data} \end{array} \right) \begin{array}{c} \xrightarrow{\mathbb{H}} \\ \xleftarrow{\mathbb{B}} \end{array} \left(\begin{array}{c} \text{Algebraic} \\ \text{Data} \end{array} \right)$$

such that:

Criterion 1.1 \mathbb{H} is homotopically defined.

Criterion 1.2 $\mathbb{H}\mathbb{B}$ is naturally equivalent to 1.

Criterion 1.3 The Topological Data has a notion of *connected*.

Criterion 1.4 For all Algebraic Data A , the Topological Data $\mathbb{B}A$ is connected.

Criterion 1.5 If the topological Data X has underlying space X which is the union of a family $\mathcal{U} = \{U_\lambda, \lambda \in \Lambda\}$ of open sets, and each corresponding Topological Data U_ν is connected for all finite intersections U_ν of the sets of \mathcal{U} , then

(a) X is connected, and (b) the following morphism induced by inclusions $U_\nu \rightarrow X$

$$\text{colim}_\nu \mathbb{H}U_\nu \rightarrow \mathbb{H}X$$

is an isomorphism.

We will give detail on the notion of “connected” in specific examples later: this condition does of course give a limitation on the applicability of such theorems, limitations which are familiar in algebraic topology. In fact considerable effort by many writers has gone into developing connectivity results, such as the theorems of Blakers-Massey, and of n -ad connectivity, sometimes proved by subtle homotopy theoretic arguments, or general position and transversality, see for example [MV15]. In practice, though, we would like an algebraic calculation, and these usually imply connectivity results, since often a kind of tensor product is involved, and this is trivial if one of the factors is trivial. The aim is precise algebraic colimit calculations of some homotopy types.

2 Broad and Narrow Algebraic Data

In dimension 1 the Topological Data we use is pairs (X, C) of spaces X with a set C of base points, and the Algebraic Data is groupoids, with the functor \mathbb{H} sending (X, C) to the groupoid $\pi_1(X, C) = \pi_1(X, X \cap C)$. This notation is a ruse to allow one to write $\pi_1(U, C)$ if, for example, U is a subspace of X .

For modelling homotopy type in dimensions > 1 this method has to be refined owing to the increased variety of convex sets in these dimensions. We then use at least two types of algebraic data for a model; their advantage is that while they are chosen to give equivalent categories, these categories serve different purposes. Thus in the following diagram, \mathbb{H} of Criterion 1.1 has split into ϱ and Π . Further U is the forgetful functor, and $B = U\mathbb{B}$ is the “classifying space” of the Algebraic Data, whose definition usually involves the Broad Algebraic Data.




Figure 1: Broad and Narrow Algebraic Data

The “Broad” Algebraic Data has a geometric type of axioms, is expressive, and useful for conjecturing and proving theorems, particularly colimit theorems, and for constructing classifying spaces. Examples given later will be double groupoids and cubical ω -groupoids.

The “Narrow” algebraic data has complicated algebraic axioms, but relates to classical theory, is useful for explicit calculation, and colimit examples have led to new algebraic constructions. Examples given later will be crossed modules and crossed complexes.

The necessarily *algebraic* proof of equivalence of these two structures allows one to get the best of both worlds. The hardest part of the equivalence is usually the natural equivalence $\lambda\gamma \simeq 1$, which shows that the broad data can be recovered from the narrow data. Also the functor γ usually involves choices. The whole relationship is a key to the power of the theory, and is important in developing aspects of it. Of course the use of various geometric models such as discs, simplices, cubes, for defining invariants and constructions is standard in homotopy theory.

We will give more details in Section 12 of the case when the Topological Data is that of Filtered Spaces. The Narrow Algebraic Data is then crossed complexes, which date back in essence to Blakers, [Bla48], and were developed by Whitehead, [Whi49]. The Broad Algebraic Data is that of cubical ω -groupoids, [BH81].

Most other aspects are explained in [BHS11], to which this article gives an introduction.

Another example which we plan to explain in Part II is the work in [BL87, BL87a], where the

Topological Data is that of n -cubes of pointed spaces, the Broad Algebraic Data is that of cat- n -groups, later called cat ^{n} -groups, and the Narrow Algebraic Data is that of crossed n -cubes of groups, [ES87].

Many categories \mathcal{AD} of Algebraic Data, either broad or narrow, have in our examples also a family of subcategories \mathcal{AD}^n classified by dimension $n \geq 0$. and these come with forgetful functors $\Phi^n : \mathcal{AD}^n \rightarrow \mathcal{AD}^{n-1}$. It usually turns out that Φ^n is a bifibration of categories, i.e. a fibration and an opfibration (or cofibration), and even has left and right adjoints. As spelled out in [BHS11, Appendix B], this helps the calculation of colimits in \mathcal{AD}^n using their determination in \mathcal{AD}^{n-1} . For example, in order to calculate colimits of groupoids we build them from knowing colimits in the category of sets, i.e. the object sets of the groupoids.

The Criteria of Section 1 rule out on various grounds a number of other types of algebraic data and models, such as simplicial groups, weak ∞ -groupoids. See also the Introduction to [EI93].

3 Five Anomalies in Algebraic Topology

It is useful to consider a subject in terms of **anomalies**, i.e. aspects which do not seem to fit with the general tenor of the subject.

Anomaly 3.1 The fundamental group is nonabelian, but homology and higher homotopy groups are abelian³. □

Anomaly 3.2 The traditional Seifert-van Kampen Theorem is a theorem about groups, but does not compute the fundamental group of the circle, which is THE basic example in algebraic topology. □

Anomaly 3.3 Traditional algebraic topology is fine with composing paths but does not allow for the **algebraic expression** of




From left to right gives **subdivision**. From right to left should give **composition**. What we need for higher dimensional, nonabelian, local-to-global problems is:

Algebraic Inverses to Subdivision.

□

³The early workers in algebraic topology were hoping for higher dimensional nonabelian versions of the fundamental group, and so Čech's paper on higher homotopy groups presented to the 1932 ICM at Zurich was not welcomed, and it was even felt they could not be different from the homology groups. So only a small paragraph on this work appeared in the Proceedings of the ICM. See also the Obituary Notice [AF61].

Anomaly 3.4 In traditional theory, for the Klein Bottle diagram



we have to write $\partial\sigma = 2b$, not

$$\partial(\sigma) = a + b - a + b.$$

One can get more refined by working in the operator chains of the universal cover which gives one (Whitehead, Fox, see [CF63]):

$$\partial\sigma = a^{b-a+b} + b^{-a+b} - a^b + b.$$

But this is clearly more complicated, and, for subtler reasons, less precise, than the earlier formula⁴. \square

Anomaly 3.5 The product $\Delta^n \times \Delta^1$ of a simplex with the 1-simplex, i.e. a unit interval, is not a simplex, although it has a simplicial subdivision, and this leads to awkwardness in dealing with homotopies in simplicial theory. \square

Anomaly 3.6 There is no easily defined **multiple composition** of simplices. \square

All the Anomalies 3.1 –3.6 can be resolved by using in some way groupoids and their cubical developments. While group objects in groups are just abelian groups, group objects in groupoids are equivalent to J.H.C. Whitehead's crossed modules, see Definition 10.1, which he used to describe the structure of

$$\pi_2(X, A, c) \rightarrow \pi_1(A, c),$$

a major example of a nonabelian structure in higher homotopy theory. This gives a functor

$$\Pi : (\text{pointed pairs}) \rightarrow (\text{crossed modules}).$$

How to compute it? We could do with a Seifert-van Kampen type theorem!

4 The origins of algebraic topology

The early workers in algebraic topology, before the subject had that name, see [Die89], wanted to define numerical invariants using cycles modulo boundaries but were not too clear about what these

⁴Actually the last formula takes place in the cellular chains of the universal cover, and dealing with modules over the group ring of a fundamental group has advantages, as shown in [CF63]. Nonetheless, information is lost, as shown in [BHS11], and mentioned in [Whi49].

were! Then Poincaré introduced formal sums of oriented simplices and so the possibility of the equation $\partial\partial = 0$, so that every boundary was a cycle. For more on the change from oriented to ordered homology, see [Bar95].

The idea of formal sums of domains came from an area with which many were concerned, see articles in [Jam], namely integration theory, where one could write “formally”

$$\int_C f + \int_D f = \int_{C+D} f.$$

This link to integration theory automatically gives an abelian theory.

There was also a strong link to group theory through the study of Riemann surfaces.

In our account we use homotopically defined functors which involve actual compositions for the algebraic operations. The use of these in sufficient generality took much later developments in category theory, which itself led to familiarity with partial algebraic operations. The aim was to find and use algebraic structures which better model the geometry, and in particular the interactions of spaces.

5 Enter groupoids

I was led into this area in the 1960s through writing a text on topology⁵, and in particular on the fundamental group $\pi_1(X, c)$ of a space X with base point c . The Seifert-van Kampen Theorem determines the fundamental group of a union of connected, pointed spaces. Thus the square of groups and morphisms induced by inclusions

$$\begin{array}{ccc} \pi_1(U \cap V, c) & \longrightarrow & \pi_1(V, c) \\ \downarrow & & \downarrow \\ \pi_1(U, c) & \longrightarrow & \pi_1(X, c) \end{array} \tag{1}$$

is a pushout of groups if U, V are open and $c \in U \cap V$, which is path connected. A proof by verifying the universal property, and more generally for unions of many sets, was given by Crowell in [Cro59], following lectures by Fox. Note that the proof by the universal property requires neither a previous proof that the category of groups admits pushouts or colimits, nor a specific construction of them.

If $U \cap V$ is not connected, there is no reasonable choice of base point, and of course a standard example of this is the circle S^1 . A sensible resolution is to hedge your bets, not be tethered to a single point, but use a set of base points appropriate to the geometry. Examples of such choices are given in the following pictures, where the second depicts a union of three sets:

The right hand figure shows where a choice of one base point would destroy any symmetry in the situation; this problem is forcefully made in the paper on applications in partial differential equations,

⁵This was published as [Bro68], and is currently available as [Bro06].




Figure 2: Two unions of non connected spaces

[PR15]. The left hand figure is used in [ACB15] to illustrate a failure of both of two forms of the Phragmen-Brouwer Property. The following Theorem was published in [Bro67], and developed in the 1968, 1988, differently titled, editions of [Bro06]. It has also been presented in [Hig71, Zi72].

Theorem 5.1 *Let the space X be the union of open sets U, V with intersection W . The fundamental groupoid $\pi_1(X, C)$ on a set C of base points and the morphisms induced by inclusion gives a pushout of groupoids*

$$\begin{array}{ccc} \pi_1(W, C) & \longrightarrow & \pi_1(V, C) \\ \downarrow & & \downarrow \\ \pi_1(U, C) & \longrightarrow & \pi_1(X, C) \end{array}$$

if C meets each path component of U, V, W . □

The paper [BRS84] considers a space X with a cover by open sets $U_\lambda, \lambda \in \Lambda$ and a set C of “base points” which meets each path component of each 1, 2, 3 -fold intersection of sets of the cover. Then $\pi_1(X, C)$ is given by a coequaliser of groupoids of the form

$$\bigsqcup_{(\mu, \nu) \in \Lambda^2} \pi_1(U_\mu \cap U_\nu, C) \rightrightarrows \bigsqcup_{\lambda \in \Lambda} \pi_1(U_\lambda, C) \xrightarrow{c} \pi_1(X, C). \quad (2)$$

The proofs of these theorems are direct generalisation of that in [Cro59], i.e. by verifying the universal property: groupoids are in some sense optimal for this theorem. The proof does not rely on knowing that pushouts exist for groupoids, nor how to compute them.

The theorem gives the transition from topology to algebra. The surprise to me of this kind of result is that any particular group $\pi_1(X, c)$ is completely determined by the theorem, but as part of a much bigger structure which is in essence a model of the homotopy 1-type. So we have a process:

$$(X, C) = (\text{union}) \xrightarrow{\text{SvK Theorem}} \pi_1(X, C) \xrightarrow{(\text{combinatorics})} \pi_1(X, c). \quad (3)$$

However it may be that we do not want to reduce to some $\pi_1(X, c)$: for example a choice of a $c \in C$ may obscure any symmetry of the situation.

One reason for the surprise is that exact sequences are a firm part of algebraic topology: for adjacent dimensions they usually give information only up to extension, or even less. That was the situation in the work with a Mayer-Vietoris sequence in nonabelian cohomology in [Bro65].

A possible reason for the success seemed to be that groupoids have structure in dimensions both 0 and 1, and this is surely needed to model homotopy 1-types. Philip Higgins told me of a dictum of Philip Hall: “You should seek an algebra which models the geometry and not try to force the geometry into a particular algebraic mode simply because it is more familiar.” Eldon Dyer had a dictum that you should carry around the maximum structure as long as possible. In particular, throwing away the groupoid structure to get to a group is likely to be less productive. He also suggested that I should take a hard line in pointing out that a connected space might be the union of two open sets each with 50 path components and the number of path components of the intersection being, say, 546.

Grothendieck wrote in [Gro, 24 April, 1983]:

“... the choice of a base point, and the 0-connectedness assumption, however innocuous they may seem at first sight, seem to me of a very essential nature. To make an analogy, it would be just impossible to work at ease with algebraic varieties, say, if sticking from the outset (as had been customary for a long time) to varieties which are supposed to be connected. Fixing one point, in this respect (which wouldn’t have occurred in the context of algebraic geometry) looks still worse, as far as limiting elbow-freedom goes! Also, expressing a pointed 0-connected homotopy type in terms of a group object mimicking the loop space (which isn’t a group object strictly speaking), or conversely, interpreting the group object in terms of a pointed “classifying space”, is a very inspiring magic indeed what makes it so inspiring it that it relates objects which are definitely of a very different nature lets say, “spaces” and “spaces with group law”. The magic shouldn’t make us forget though in the end that the objects thus related are of different nature, and cannot be confused without causing serious trouble.”

All this suggests that for higher dimensions we should look for groupoid like algebraic systems with structure in dimensions $0, 1, \dots, n$. This was one of the exciting possibilities I considered in 1965, when it seemed that aspects of the proof of Theorem 5.1 could generalise to the next dimension, see Section 6.

To model gluing you need to model spaces and maps. But some of the maps may involve identifications in dimension 0, i.e. of points. This can be handled in the algebra of groupoids, see [Hig71, Bro06], but clearly cannot even be formulated in an account which allows only path connected spaces with one base point.

A basic example for such identification of points is to consider the map of pairs of spaces

$$p : ([0, 1], \{0, 1\}) \rightarrow (S^1, \{1\}),$$

which just identifies 0 and 1. On taking π_1 we get the universal morphism of groupoids

$$\mathcal{I} \rightarrow \mathbb{Z},$$

where \mathcal{I} is the groupoid with two objects $0, 1$ and exactly one arrow $\iota : 0 \rightarrow 1$. The groupoid \mathcal{I} is a generator for the category of groupoids, as the infinite cyclic group \mathbb{Z} is for the category of groups.

To see how this fits with the overall philosophy of Section 1, we take the Topological Data to be a space X and a set C ; the Algebraic Data is groupoids; the functor \mathbb{H} gives $\pi_1(X, C)$; the functor \mathbb{B} assigns to a groupoid G the classifying space $B(G)$ together with the set $Ob(G)$. Thus Anomaly 3.2 is resolved by the use of the fundamental groupoid on a set of base points.

See also [Bro87] for wider uses of groupoids.

The next Section explains how the use of groupoids also suggested the possibility of resolving Anomaly 3.1.

6 Higher dimensions?

The abelian nature of higher homotopy groups can be explained as follows: the homotopy classes of maps $(I^n, \partial I^n) \rightarrow (X, x)$ allows for $n > 1$ the structure of a set with more than 1 compatible group structures, and a simple argument then shows that any two of these group structures coincide and are abelian.

However it was known to the Grothendieck school in the 1960s, and was published independently in [BS76b], that a set with the two compatible structures of group and groupoid is not necessarily abelian, and is equivalent to the structure known as a crossed module, as defined in Definition 10.1. We see later the occurrence of these structures in homotopy theory through second relative homotopy groups; they also occurred in homological algebra, for representing elements of $H^3(G, M)$ where G is a group and M is a G -module.

So the question could be formulated: are there higher homotopical invariants with structure in dimensions from 0 to n ⁶? Are there colimit theorems in higher homotopy theory? If so, what would they calculate?

The proof of the groupoid theorem seemed to generalise to dimension 2, at least, if one had the right algebra of double groupoids, and the right gadget, a strict homotopy double groupoid of a space. So this was an “idea for a proof in search of a theorem”. Fortunately I found the book [Ehr65]: it contained the notion of double category and groupoid, which seemed the right concepts for resolving Anomaly 3.3.

First we describe briefly the basic algebra of double groupoids, due to Charles Ehresmann.

We think of a square in a space X as a map $I^2 \rightarrow X$. Such a square is thought of as having four “edges”, each a path in X , as shown in the following picture.

⁶The research method adopted was to look for obstructions to this prospect. If they were found, that could be interesting. If they were, one by one, overcome, that would be even more interesting!

$$\begin{array}{ccc}
& \xrightarrow{\partial_1^-} & \\
\partial_2^- \downarrow & \square & \downarrow \partial_2^+ \\
& \xrightarrow{\partial_1^+} &
\end{array}
\quad
\begin{array}{c}
\rightarrow 2 \\
\downarrow 1
\end{array}
\tag{4}$$

Analogously, in a double groupoid G we have objects called squares and each square α has “edges” ∂_i^\pm for $i = 1, 2$, which lie in groupoids G_1, G_2 , both with the same set of objects G_0 . One makes some obvious geometric conditions, so that a square has just 4 vertices.

Compositions in a double groupoid are allowed as follows: squares α, β have a composite $\alpha +_1 \beta$ if and only if $\partial_1^+ \alpha = \partial_1^- \beta$, and the composition $+_1$ gives a groupoid structure. Similarly we have a composition $+_2$.

$$\begin{array}{c}
\boxed{\alpha} \\
\boxed{\beta}
\end{array}
= \begin{bmatrix} \alpha \\ \beta \end{bmatrix} = \alpha +_1 \beta$$

$$\begin{array}{|c|c|}
\hline
\alpha & \gamma \\
\hline
\end{array}
= [\alpha \quad \gamma] = \alpha +_2 \gamma$$

$$\begin{array}{|c|c|}
\hline
\alpha & \gamma \\
\hline
\beta & \delta \\
\hline
\end{array}
= \begin{bmatrix} \alpha & \gamma \\ \beta & \delta \end{bmatrix}
\quad
\begin{array}{c}
\rightarrow 2 \\
\downarrow 1
\end{array}$$

That each is to be a morphism for the other gives the interchange law:

$$(\alpha \circ_2 \gamma) \circ_1 (\beta \circ_2 \delta) = (\alpha \circ_1 \beta) \circ_2 (\gamma \circ_1 \delta)
\tag{5}$$

for which the necessary and sufficient condition is that all the compositions in the equation are defined. This equation illustrates that a 2-dimensional picture can be more comprehensible than a 1-dimensional equation.

In these double groupoids the horizontal edges and vertical edges may come from different groupoids.

Note also that multiple compositions can be constructed by considering “compatible” arrays (a_{ij}) of elements of a double groupoid, and their composite $[a_{ij}]$. Compatible means that each a_{ij} is composable with its neighbours. This easy expression of “algebraic inverses to subdivision” is one of the chief advantages of the cubical approach. It allows for the potential of resolving Anomaly 3.3, and also Anomaly 3.1, since we will see later that double groupoids can be much more complicated than abelian groups. Indeed, classification of the most general double groupoids is not well understood.

7 Need also “Commutative cubes”

In dimension 1, we still need the 2-dimensional notion of commutative square:

$$\begin{array}{c} \xrightarrow{c} \\ a \downarrow \quad \downarrow d \\ \xrightarrow{b} \end{array} \quad ab = cd, a = cdb^{-1} \quad (6)$$

An easy result is that any composition of commutative squares is commutative. In ordinary equations:

$$ab = cd, ef = bq \implies aef = abq = cdq.$$


The commutative squares in a groupoid G form a double groupoid $\square G$.

This fact is a key part of one proof of Theorem 5.1, to verify that a morphism to a groupoid is well defined. The essence is as follows: consider a diagram in a groupoid

$$\begin{array}{|c|c|c|c|c|} \hline & b & & & \\ \hline 1 & & & & 1 \\ \hline & & & & \\ \hline & & & & \\ \hline & & & & \\ \hline & a & & & \\ \hline \end{array} \quad (7)$$

Suppose each individual square is commutative, and the two vertical outside edges are identities. Then we easily deduce $a = b$.

For the next dimension we therefore expect to need to know what is a commutative cube, and this is expected to be in a double groupoid in which horizontal and vertical edges come from the same groupoid: $G_1 = G_2$.



We want the “composed faces” to commute! What can this mean?

We might say the top face is the composite of the other faces: so fold them flat to give the left hand diagram of Fig. 3, where the dotted lines show adjacent edges of a “cut”. We indicate how to glue these edges back together by extra squares which are a different kind of “degeneracy” in the right hand diagram of this Figure.




Figure 3: “Composing” five faces of a cube

Thus if we write the standard double groupoid identities in dimension 2 by

$$\square = ||$$

where a solid line indicates a constant edge, then the new types of square with commutative boundaries are written

$$\ulcorner \urcorner \llcorner \lrcorner$$

These new kinds of “degeneracies” are called **connections**, [BS76b]⁷. The laws on connections are:

$$[\ulcorner \lrcorner] = || \quad \left[\begin{array}{c} \ulcorner \\ \lrcorner \end{array} \right] = = \quad (\text{cancellation})$$

$$\left[\begin{array}{cc} \ulcorner & = \\ || & \ulcorner \end{array} \right] = \ulcorner \quad \left[\begin{array}{cc} \lrcorner & || \\ = & \lrcorner \end{array} \right] = \lrcorner \quad (\text{transport})$$

These are equations on turning left or right, and so are a part of “2-dimensional algebra”.

The terms **transport law** and **connection** came from laws on path connections in differential geometry, see [BS76b].

Any well defined 2-dimensional composition of identities and connections is called **thin**. A thin square is entirely determined by its boundary.

Full details are given in [BHS11, Section 6.7]. This proof uses “2-dimensional rewriting”, as does the proof for strict cubical ω -groupoids in Chapter 13 of [BHS11]. The account for the case of strict cubical ω -categories in [Hig05] uses “3-dimensional rewriting”.

Double groupoids allow for: multiple compositions, 2-dimensional formulae, and 2-dimensional rewriting. As an example, we get a “rotation”

$$\sigma(\alpha) = \left[\begin{array}{cc} || & \ulcorner \\ \lrcorner & \alpha \\ = & \lrcorner \end{array} \right]$$

⁷The latex code which was developed for this notation for the identities and connections and used for example in [BHS11] is available at the following <http://www.groupoids.org.uk/connections-texcode.html>, Its advantages can be seen for interpreting diagram (9). The notation was first published in [Bro82].

As a taster to “2-dimensional rewriting” we note that making two interpretations of the following diagram

$$\left[\begin{array}{c|c|c|c|c} || & \ulcorner & = & = & = \\ \hline || & || & \square & \square & \square \\ \hline \llcorner & \alpha & \urcorner & \square & \square \\ \hline = & \lrcorner & || & \square & \square \\ \hline \square & \square & || & \ulcorner & = \\ \hline \square & \square & \llcorner & \beta & \urcorner \\ \hline \square & \square & \square & || & || \\ \hline = & = & = & \lrcorner & || \end{array} \right] . \quad (9)$$

shows that $\sigma \begin{bmatrix} \alpha & \beta \end{bmatrix} = \begin{bmatrix} \sigma\alpha \\ \sigma\beta \end{bmatrix}$. See [BHS11, Theorem 6.4.10]. Interpreting this proof in a homotopy double groupoid shows in principle how to construct explicit homotopies. Diagrammatic arguments using the rules for connections are a central feature of that book.

It turns out that connections are what is needed, [BS76a], to obtain that double groupoids with connections are equivalent to crossed modules; these two structures give respectively broad and narrow algebraic data.

This result was found by trying to construct examples of double groupoids. The first example, given by Ehresmann, [Ehr65], was that of commutative squares in a group or groupoid G . The next example, say in the case G is a group, was to look at squares which commuted up to an element of a subgroup N , say: to get compositions one quickly found N had to be normal in G .

Finally, we looked at a morphism $\nu : N \rightarrow G$ of groups and quintuples $(n; a, b, c, d)$ where $n \in N, a, b, c, d \in G$ and $\nu n = b^{-1}a^{-1}cd$. Compositions are easy to define if G operates on N satisfying CM1) of Definition 10.1, but to get the interchange law we need exactly also CM2). The thin squares are of the form $(1; a, b, c, d)$. For more details, see also [BM99].

These ideas led to the notion of “higher dimensional group theory” as a theory to be developed in the spirit of group theory, but using algebraic structures involving partial operations whose domains were defined by geometric conditions. This scheme fitted with the work of Higgins in [Hig63].

8 Cubical sets in algebraic topology

Dan Kan’s first paper, [Kan55], was cubical, relying clearly on geometry and intuition. It was then found that cubical groups, unlike simplicial groups, were not Kan complexes, and there was also a problem on realisation of cartesian products. The simplicial methods seemed entirely adequate, and the cubical theory seemed unfixable; so cubical methods were generally abandoned for the simplicial, although many workers found them intuitive and useful in certain contexts. The book [Mas80] gives a cubical approach to singular homology.

The work with Chris Spencer on double groupoids in the early 1970s, [BS76a], found it necessary to introduce an extra and new kind of “degeneracy” in cubical sets, using the monoid structures

$$\max, \min : [0, 1]^2 \rightarrow [0, 1].$$

We called these “connection operators”. The axioms for these in all dimensions were published in [BH81]. Andy Tonks proved in [Ton92] that cubical groups with connections are Kan complexes! G. Maltsiniotis in [Mal09] showed that up to homotopy, connections correct the realisation problem. Patchkoria in [Pat12] gave a cubical approach to derived functors. Cubical sets have analogously been used for work on motives, see [Vez14], and the references there⁸.

Cubical sets have also been used in concurrency theory, see [Gou03]: if many computers run databases, then this can be envisaged as a system with many-dimensional time. They are also used in Homotopy Type Theory, [CCHM]. A recent work, [RZ16], relates cubical sets with connections to the simplicial method of necklaces.

Independently of these facts, we deal with cubical sets with connections and compositions, where the compositions have properties analogous to those of paths in a space; if all the compositions are groupoid structures, we get a cubical ω -groupoid, [BH81], [BHS11].

9 Strict homotopy double groupoids?

In 1965, inspired by a definition in [Ehr65] which seemed to give a concept appropriate to resolve Anomaly 3.3, I started to seek a strict homotopy double groupoid of a space. Group objects in groups are abelian groups, by the interchange law. Chris Spencer and I found out in the early 1970s that group objects in groupoids are more complicated, in fact equivalent to Henry Whitehead’s crossed

⁸Vezzani has explained to me the use as follows: The cubical theory was better suited than the simplicial theory when dealing with (motives of) perfectoid spaces in characteristic 0. For example: degeneracy maps of the simplicial complex Δ in algebraic geometry are defined by sending one coordinate x_i to the sum of two coordinates $y_j + y_{j+1}$. When one considers the perfectoid algebras obtained by taking all p -th roots of the coordinates, such maps are no longer defined, as $y_j + y_{j+1}$ doesn’t have p -th roots in general. The cubical complex, on the contrary, is easily generalized to the perfectoid world.

modules, see Definition 10.1, as was known earlier to some, see [BS76b]. The important point was that double groupoids could be highly non trivial⁹.

In the early 1970s Chris Spencer, Philip Higgins and I developed a lot of understanding of:

- (i) relations between double groupoids and crossed modules; and
- (ii) algebraic constructions on the latter, including induced crossed modules, and colimit calculations.

In June, 1974, Phil and I, stimulated by the last week of his supported visit to Bangor, argued as follows:

Strategic Analysis 9.1

1. Whitehead had a subtle theorem (1941–1949), [Whi41, Whi49], that

$$\pi_2(A \cup \{e_\lambda^2\}, A, c) \rightarrow \pi_1(A, c)$$

is a free crossed module. So this was one, indeed seemingly the only, example of a universal property in 2-dimensional homotopy theory.

2. Thus Whitehead's theorem should be a consequence of our conjectured theorem, if that so far unformulated theorem was to be any good.
3. But Whitehead's theorem was about second *relative* homotopy groups.
4. So we should look for a homotopy double groupoid in a relative situation, (X, A, c) .
5. So we tried what seemed the simplest idea, i.e. to look at maps of the square $I^2 \rightarrow X$ which took the edges of the square to A and the vertices to c , and consider homotopy classes of these.
6. Because of all the preliminary work with Chris and Phil, this worked like a dream! It was submitted 1975, and published severely cut as [BH78].

□

10 Crossed Modules

Definition 10.1 A *crossed module* consists of a morphism of groupoids $\mu : M \rightarrow P$ of groupoids such that μ is the identity on objects, M is totally disconnected, and there is also given an action of P on M written $(p, m) \mapsto {}^p m$, where if $p : x \rightarrow y$ and $m \in M(x)$, then ${}^p m \in M(y)$. The usual rules for such an action should be satisfied as well as for all $m, n \in M, p \in P$

CM1) $\mu({}^p m) = p(\mu m)p^{-1}$, and

CM2)¹⁰ $mnm^{-1} = {}^{\mu m} n$.

Crossed modules and their morphisms form a category XMod.

□

⁹The further question was: could the leap to these higher dimensional versions of groups be analogous to the leap from real analysis to many dimensional analysis?

¹⁰To my knowledge, this rule was first stated for the group case in a footnote on p.422 of [Whi41].


If P is a group, the notion of free crossed P -module $C(R) \rightarrow P$ on a function $w : R \rightarrow P$ can be expressed by the pushout of crossed modules:

$$\begin{array}{ccc} (1 \rightarrow F(R)) & \longrightarrow & (1 \rightarrow P) \\ \downarrow & & \downarrow \\ (F(R) \xrightarrow{1} F(R)) & \longrightarrow & (C(R) \rightarrow P), \end{array} \quad (10)$$


where $F(R) \rightarrow F(R)$ is the identity crossed module. Whitehead's Theorem solves the “Klein Bottle Anomaly” [3.4](#): the element σ is the generator of $\pi_2(K^2, K^1, x)$ as a free crossed $\pi_1(K^1, x)$ -module. This crossed module gives “nonabelian chains” in dimension ≤ 2 . Whitehead's proof^{[11](#)} of his theorem uses transversality and knot theory, and seems unlikely to suggest generalisations.

11 Groupoids in higher homotopy theory?

Consider second relative homotopy groups $\pi_2(X, A, c)$.




where double lines show constant paths. Note that the definition involves choices, and is unsymmetrical with respect to directions. This is unaesthetic! Further, all compositions are on a line:



All this is necessary to obtain a group structure, and, by tradition, the structure has to be a group!

In 1974 Philip Higgins and the author considered a symmetrical version of relative homotopy groups, namely $\varrho_2(X, A, C)$, the homotopy classes rel vertices of maps $[0, 1]^2 \rightarrow X$ with edges mapped into A and vertices into C :



¹¹A modern and selfcontained exposition of his proof is in [\[Bro80\]](#).


$$\varrho_2(X, A, C) \rightrightarrows \pi_1(A, C) \rightrightarrows C$$

This allowed for the childish idea of gluing two squares if the right side of one is the same as the left side of the other. This gives a partial algebraic operation defined under geometric condition.

There is a horizontal composition

$$\langle\langle\alpha\rangle\rangle +_2 \langle\langle\beta\rangle\rangle = \langle\langle\alpha +_2 h +_2 \beta\rangle\rangle$$

of classes in $\varrho_2(X, A, C)$, where thick lines show constant paths.



Intuition: gluing squares exactly is a bit too rigid; varying edges in X is too free; but “varying edges in A ” seems just about right.

To show $+_2$ well defined, let $\phi : \alpha \equiv \alpha'$ and $\psi : \beta \equiv \beta'$, and let $\alpha' +_2 h' +_2 \beta'$ be defined. We get Figure 4 in which thick lines denote constant paths. Can you see why the ‘hole’ can be filled appropriately?




Figure 4: Showing $+_2$ is well defined on ϱ

We find that $\varrho(X, A, C)$ has in dimension 2 compositions in directions 1 and 2 satisfying the interchange law and is a double groupoid with connections, containing as an equivalent substructure the classical

$$\Pi(X.A, C) = (\pi_2(X.A, C) \rightarrow \pi_1(A, C)),$$

a crossed module over a groupoid. All this needs proof, but in this dimension is not too hard. Thus ϱ gives in this case the “broad” structure and Π the “narrow” structure. Now we can directly generalise the 1-dimensional proof: since one has a homotopy double groupoid which is “equivalent” to crossed modules but which can express:

- algebraic inverse to subdivision
- the notion of commutative cubes, with the property that any multiple composition of commutative cubes is commutative.

A key deformation idea is shown in Figure 5 We need to deform the bottom subdivided square into




Figure 5: Partial deformation of a subdivided square

a subdivided square for which all of its subsquares define an element of $\varrho(X, A, C)$. This explains the connectivity assumptions for the theorem. It is to express this diagram that $\varrho(X, A, C)$ is designed.

We end up with a 2-d SvK theorem, namely a pushout of crossed modules:

$$\begin{array}{ccc}
 \Pi(X, A, C) \cap U \cap V & \longrightarrow & \Pi(X, A, C) \cap V \\
 \downarrow & & \downarrow \\
 \Pi(X, A, C) \cap U & \longrightarrow & \Pi(X, A, C)
 \end{array} \tag{11}$$

if $X = U \cup V$, U, V open, and the intersections of (X, A, C) with $U, V, U \cap V$ are connected in the following sense:

Definition 11.1 We say (X, A, C) is *connected* if

- (i) $\pi_0(C) \rightarrow \pi_0(A), \pi_0(C) \rightarrow \pi_0(X)$ are surjective; and
- (ii) any map $(I, \partial I) \rightarrow (X, C)$ is deformable into A rel end points.

□

The huge generalisation of Whitehead's theorem given by the pushout of diagram (11), says that you glue homotopy 2-types, by gluing crossed modules as a pushout. A consequence is the following pushout, for A, X connected and $f : A \rightarrow X$:

$$\begin{array}{ccc}
 (1 \rightarrow \pi_1(A, a)) & \longrightarrow & (1 \rightarrow \pi_1(X, f(a))) \\
 \downarrow & & \downarrow \\
 (\pi_1(A, a) \xrightarrow{1} \pi_1(A, a)) & \longrightarrow & (\pi_2(X \cup_f C(A), X, f(a)) \rightarrow \pi_1(X, f(a))).
 \end{array} \tag{12}$$

Whitehead's Theorem is the case A is a wedge of circles.

The results enable some nonabelian computations of homotopy 2-types, see [BHS11, Chapters 4,5]. There are methods to determine explicitly the module π_2 from this algebraic model of the 2-type, but that module is a pale shadow of the 2-type.

The proof of all this works first in the category of double groupoids with connection, and then uses the equivalence with crossed modules given in [BS76a]. The notion of connection is the precisely necessary tool for both parts of the proof.

To define the functor \mathbb{B} on crossed modules it is best to generalise to crossed complexes and their relation with filtered spaces.

12 Filtered spaces

The success of the 2-d work led to a search for the n -dimensional idea, and in view of the work of [Bla48] and [Whi49] it was natural to look at filtered spaces:

$$X_* := X_0 \subseteq X_1 \subseteq \cdots \subseteq X_n \subseteq \cdots \subseteq X \quad (13)$$

With the obvious morphisms, these form a category \mathbf{FTop} . Examples of filtered spaces are:

- (i) the skeletal filtration of a CW-complex: e.g. Δ_*^n, I_*^n .
- (ii) $\bullet \subseteq A \subseteq \cdots \subseteq A \subseteq X \subseteq X \subseteq \cdots$, where the first X occurs at position n ;
- (iii) for a pointed topological space X , the free monoid $FM(X)$ on X may be filtered by word length, cf, [BaBr93].

Now we need to define the notion of a cubical set with compositions and connections and then check that for a filtered space X_* we get an example of this if we define $R(X_*)$ by

$$R_n(X_*) = \mathbf{FTop}(I_*^n, X_*).$$

Of course the compositions here do not form groupoids.

Now define a *thin* homotopy $h : a \equiv b$ between elements $a, b \in R_n(X_*)$ to be a homotopy through filtered maps and rel vertices of the cube. Such a homotopy is a particular kind of element of $R_{n+1}(X_*)$. The thin homotopy class of an element $a \in R_n(X_*)$ is written $[a]$.

Theorem 12.1 *The thin homotopy classes of elements of $R(X_*)$ form a cubical set with connections written $\varrho(X_*)$ with projection $p : R(X_*) \rightarrow \varrho(X_*)$. Then:*

- (i) *the compositions on $R(X_*)$ are inherited to give $\varrho(X_*)$ the structure of strict cubical ω -groupoid;*
- (ii) *$\varrho(X_*)$ is a Kan complex in which every box has a unique thin filler, where an element $\alpha = [a] \in \varrho_n(X_*)$ is thin if it has a representative a such that $a(I^n) \subseteq X_{n-1}$;*
- (iii) *the projection p is a Kan fibration of cubical sets.* □

The proof was given in [BH81a], and full details are in [BHS11]; that of (i) uses cubical versions of Whitehead's notions of expansions and collapsing, and it seems significant that there is exactly enough "filtered room" to make the proof work.

The proof that $R(X_*) \rightarrow \varrho(X_*)$ is a fibration, relies on a nice use of geometric cubical methods, and the Kan condition. One applies the Kan condition by modelling in subcomplexes of real cubes and using expansions and collapsings of these.

A consequence of the fibration property is the following, which we give here because it emphasises the use of multiple compositions:

Corollary 12.2 (Lifting composable arrays) *Let $(\alpha_{(i)})$ be a composable array of elements of $\varrho_n(X_*)$. Then there is a composable array $(a_{(i)})$ of elements of $R_n(X_*)$ such that for all (i) , $p(a_{(i)}) = \alpha_{(i)}$. \square*

Thus the weak cubical infinity groupoid structure of $R(X_*)$ has some kind of control by the strict infinity groupoid structure of $\varrho(X_*)$.

13 Crossed complexes and their classifying space

Associated functorially with a filtered space X_* is a homotopical invariant $\Pi(X_*)$ called a *crossed complex*, due in the case X_0 is a point to Blakers, [Bla48], and Whitehead, [Whi49]. It involves the crossed module of groupoids

$$\pi_2(X_2, X_1, X_0) \rightarrow \pi_1(X_1, X_0),$$

and in dimensions > 2 the boundary map

$$\pi_n(X_n, X_{n-1}, x) \rightarrow \pi_{n-1}(X_{n-1}, X_{n-2}, x), x \in X_0$$

together with the operation of $\pi_1(X_1, X_0)$ on all these groups. So we have a functor $\Pi : \mathbf{FTop} \rightarrow \mathbf{Crs}$.

For our purposes, a key result is that the algebraic equivalence of categories

$$\gamma : \omega\text{-Gpd} \rightarrow \mathbf{Crs}$$

takes $\varrho(X_*)$ to $\Pi(X_*)$. So we may use whatever is best in a given situation. It also confirms that the axioms for crossed complexes and for ω -groupoids are the right ones. Thus the algebra matches the geometry.

Now we can define the classifying space $B(C)$ of a crossed complex C . The cubical definition fits better with other themes. So we first define the cubical set $N(C)$ in dimension n by

$$N(C)_n = \mathbf{Crs}(\Pi(I_*^n), C).$$

We then define

$$B(C) = |N(C)|,$$

the geometric realisation of the cubical set $N(C)$.

But C can be filtered by its truncations $C^{(n)}$, the crossed complex which agrees with C up to and including dimension n and above that is trivial. So we get a filtered space $\mathbb{B}(C)$ which in dimension n is $B(C^{(n)})$. The proof of connectivity of $\mathbb{B}(C)$ uses the adjointness

$$\text{Cub}(K, N(C)) \cong \text{Crs}(\Pi(|K|_*, C)).$$

We also need to discuss connectivity of a filtered space X_* . The motivation of the condition is to allow the deformations used in the proof of the Seifert-van Kampen Theorem, namely that the functor ϱ , and hence also Π , preserves certain colimits.

Definition 13.1 A filtered space X_* is *connected* if the following conditions hold: for all $n > 0$
 $(\phi 0)$: the induced function $\pi_0 X_0 \rightarrow \pi_0 X_n$ is surjective and,
 (ϕn) : $\pi_j(X_{n+1}, X_n, v) = 0$ for all $j \leq n$, and $v \in X_0$. □

There are other equivalent forms of this definition, obtained using homotopy exact sequences. It is standard that these conditions are satisfied if X_* is the skeletal filtration of a CW-complex.

Theorem 13.2 Let X_* be a filtered space and let $\mathcal{U} = \{U^\lambda, \lambda \in \Lambda\}$ be an open cover of X . For each $\nu \in \Lambda^n$ let U^ν denote the intersection of the sets U^{ν_i} and let U_*^ν be the filtered space formed by intersection with the $X_i, i \geq 0$. Suppose that for all $n \geq 1$ each $U_*^\nu, \nu \in \Lambda^n$ is a connected filtered space. Then

(Con) X_* is connected, and

(Iso) the following diagram is a coequaliser diagram of crossed complexes:

$$\bigsqcup_{\nu \in \Lambda^2} \Pi U_*^\nu \rightrightarrows_a^b \bigsqcup_{\lambda \in \Lambda} \Pi U_*^\lambda \xrightarrow{c} \Pi X_*, \quad \square$$

Here a, b are induced by the inclusions $U_{(\lambda, \mu)} \rightarrow U_\lambda, U_{(\lambda, \mu)} \rightarrow U_\mu$ respectively, and c is induced by the inclusions $U_\lambda \rightarrow X$.

This Higher Homotopy Seifert-van Kampen Theorem allows one easily to compute for example $\Pi(X_*)$ when X_* is the skeletal filtration of a CW-complex.

14 Why filtered spaces?

In the proof of Theorem 12.1, particularly the proof that compositions are inherited, there is exactly the right amount of “filtered room”. One then needs to evaluate the significance of that fact!

Grothendieck in “Esquisse d’un Programme”, [Gro84, Section 5], has attacked the dominance of the notion of topological space, which he says comes from the needs of analysis rather than geometry. He advocates some ideas of stratified spaces: filtered spaces are a step in that direction. A general

argument is that to describe or specify a space you need some kind of data, and that data has some kind of structure. So it is reasonable for the invariants to be defined using that structure.

The theory given so far gives an account of algebraic topology on the border between homology and homotopy but without the use of singular homology or simplicial approximation. Nonetheless it allows a geometric account of results such as:

- (i) the Brouwer degree theorem (the n -sphere S^n is $(n - 1)$ -connected and the homotopy classes of maps of S^n to itself are classified by an integer, called the degree of the map);
- (ii) the Relative Hurewicz theorem, which usually relates relative homotopy and homology groups, but is seen here as describing the morphism $\pi_n(X, A, x) \rightarrow \pi_n(X \cup CA, x)$ when (X, A) is $(n - 1)$ -connected;
- (iii) Whitehead's theorem (1949) referred to earlier;
- (iv) a generalisation of that theorem to describe the crossed module

$$\pi_2(X \cup_f CA, X, x) \rightarrow \pi_1(X, x)$$

as induced by the morphism $f_* : \pi_1(A, a) \rightarrow \pi_1(X, x)$ from the identity crossed module $\pi_1(A, a) \rightarrow \pi_1(A, a)$; and

- (v) a coproduct description of the crossed module $\pi_2(K \cup L, M, x) \rightarrow \pi_1(M, x)$ when $M = K \cap L$ is connected and $(K, M), (L, M)$ are 1-connected and cofibred.

The Fibration Theorem, part (iii) of Theorem 12.1, yields an associated family of homotopy exact sequences which, in the case X_* is a CW-filtration, corresponds to Whitehead's exact sequence in [Whi50], as shown in [BHS11, Theorem 14.7.9]. Thus the base $\varrho(X_*)$ of the fibration, captures the “linear” part of the space X , while $R(X_*)$ has the homotopy type of X .

The above cited book gives applications of monoidal closed categories to homotopy classification results, based on the key isomorphism

$$I_*^m \otimes I_*^n \cong I_*^{m+n}.$$

For example it is easy to define in the cubical setting a natural transformation

$$\eta : \varrho(X_*) \otimes \varrho(Y_*) \rightarrow \varrho(X_* \otimes Y_*),$$

basically as $[f] \otimes [g] \mapsto [f \otimes g]$, and then to transfer this to the crossed complex case using the equivalence of categories. This leads to the notion of “crossed differential algebra”, i.e. monoid objects with respect to \otimes in the category CrS , though the theory may work out better in the category $\omega\text{-Gpd}$.

A further advantage of the monoidal closed property of the category \mathbf{Crs} is to yield a new view of the Homotopy Addition Lemma, or Theorem: this gives an explicit formula for $\Pi(\Delta_*^n)$. This result is used in [Bla48], was first proved in [Hu53], is proved in [Whi78] inductively in conjunction with the Relative and Absolute Hurewicz Theorems. By contrast, in [BHS11, Theorem 9.9.4] it is proved in the category \mathbf{Crs} by regarding the filtered simplex Δ_*^n as a cone on Δ_*^{n-1} and using the detailed algebraic description of the tensor product of crossed complexes.

Interestingly, a kind of 2-dimensional form of the HAL plays a key role in Whitehead’s proof of his theorem on free crossed modules: see Figure 6, taken from [Bro80, p.148], which is used to




Figure 6: 2-dimensional Homotopy Addition Lemma

describe an element of $\pi_2(X \cup \{e_\lambda^2\}, X, x)$ as given by a product of elements of the form $(c_{i_k}^{\xi_k})^{\zeta_k}$.

The crossed complex model of homotopy types can be described as essentially a “linear” model: it does not even capture quadratic information such as Whitehead products. Grothendieck was led by this limitation of modelling of homotopy types to embark as one thread in “Pursuing Stacks”, [Gro83], on the notion of weak ∞ -groupoids, on which there is now a considerable body of work¹².

The linearity limitation can also be seen intuitively in the fact that crossed complexes can be written on a line, and crossed resolutions are constructed step by step in increasing dimensions.

By contrast, Loday’s construction of a “resolution” of a connected, pointed space in [Lod82], corrected in [Ste86], moves a step in one direction, then in an orthogonal direction, and so on, to end up with the Topological Data of connected n -cube of fibrations, and with Broad Algebraic Data that of \mathbf{cat}^n -groups, which are essentially $(n + 1)$ -fold groupoids in which one direction, say the last, is a group. The Narrow Algebraic Data is the crossed n -cubes of groups of [ES87]. More on this will be written in a second paper.

One aspect from this work which has been well taken up is the notion of nonabelian tensor product of groups¹³, defined in [BL87]. Nonetheless, this restriction to pointed spaces is yet another Anomaly!

¹² See <https://ncatlab.org/nlab/show/infinity-groupoid>. This literature currently makes little reference to cubical methods.

¹³I have kept up a bibliography on this area, available as www.groupoids.org.uk/nonabtens.html, with

Further, it can be expected that quadratic information can be recovered by using crossed bicomplexes, i.e. crossed complexes internal to the category \mathbf{Crs} .

The Algebraic Model of n -fold groupoids is considered in [BB12], but currently this work has not been linked with the work on higher order van Kampen theorems, i.e. following the philosophy of Section 1.

References

- [AF61] Aleksandrov, P. S. and Finikov, S. P. ‘Eduard Čech: Obituary’. *Uspehi Mat. Nauk* **16** (1 (97)) (1961) 119–126. (1 plate). Cited on p. 6
- [ACB15] Antolin-Camarena, O. and Brown, R. ‘Corrigendum to “Groupoids, the Phragmen-Brouwer Property, and the Jordan Curve Theorem” J. Homotopy and Related Structures 1 (2006) 175-183.’ *J. Homotopy and Related Structures* (2015) 10:665-672. Cited on p. 9
- [Ati02] Atiyah, M. ‘Mathematics in the 20th century’, *London Math. Soc.* **34** (2002) 1-15. Cited on p. 3
- [Bar95] Barr, M. ‘Oriented Singular Homology’. *Theory and Applications of Categories* **1** (1995) 1-9. Cited on p. 8
- [BaBr93] Baues, H. J. and Brown, R. ‘On relative homotopy groups of the product filtration, the James construction, and a formula of Hopf’. *J. Pure Appl. Algebra* **89** (1-2) (1993) 49–61. Cited on p. 21
- [BB12] Blanc, D. and Paoli, S. ‘ n -fold groupoids, n -types and n -track categories’ (2012) arXiv:1204.5101. Cited on p. 26
- [Bla48] Blakers, A. ‘Some relations between homology and homotopy groups’. *Ann. of Math.* (2) **49** (1948) 428–461. Cited on p. 5, 21, 22, 25
- [Bro65] Brown, R. ‘On a method of P. Olum’. *J. London Math. Soc.* **40** (1965) 303–304. Cited on p. 10
- [Bro67] Brown, R. ‘Groupoids and van Kampen’s theorem’. *Proc. London Math. Soc.* (3) **17** (1967) 385–401. Cited on p. 9
- [Bro68] Brown, R. *Elements of Modern Topology*. McGraw-Hill Book Co., New York (1968). Cited on p. 8, 27

currently 153 items dating from 1952. Most interest has been from group theorists.

- [Bro80] Brown, R. ‘On the second relative homotopy group of an adjunction space: an exposition of a theorem of J. H. C. Whitehead’. *J. London Math. Soc. (2)* **22** (1) (1980) 146–152. Cited on p. [18](#), [25](#)
- [Bro82] Brown, R. ‘Higher dimensional group theory’. In ‘Low-dimensional topology (Bangor, 1979)’, *London Math. Soc. Lecture Note Ser.*, Volume 48. Cambridge Univ. Press, Cambridge (1982), 215–238. Cited on p. [14](#)
- [Bro87] Brown, R. ‘From groups to groupoids: a brief survey’. *Bull. London Math. Soc.* **19** (2) (1987) 113–134. Cited on p. [11](#)
- [Bro06] Brown, R. *Topology and Groupoids*. Printed by Booksurge LLC, Charleston, S. Carolina, third edition of [[Bro68](#)]. available from amazon. (2006). Cited on p. [8](#), [9](#), [10](#)
- [BH78] Brown, R. and Higgins, P. J. ‘On the connection between the second relative homotopy groups of some related spaces’. *Proc. London Math. Soc. (3)* **36** (2) (1978) 193–212. Cited on p. [3](#), [17](#)
- [BH81] Brown, R. and Higgins, P. J. ‘On the algebra of cubes’. *J. Pure Appl. Algebra* **21** (3) (1980) 233–260. Cited on p. [5](#), [16](#)
- [BH81a] Brown, R. and Higgins, P. J. ‘Colimit theorems for relative homotopy groups’. *J. Pure Appl. Algebra* **22** (1) (1981) 11–41. Cited on p. [21](#)
- [BHS11] Brown, R., Higgins, P. J. and Sivera, R. *Nonabelian algebraic topology: filtered spaces, crossed complexes, cubical homotopy groupoids*. EMS Tracts in Mathematics Vol 15. European Mathematical Society (2011). Cited on p. [5](#), [6](#), [7](#), [14](#), [15](#), [16](#), [21](#), [24](#), [25](#)
- [BL87] Brown, R. and Loday, J.-L. ‘Van Kampen theorems for diagrams of spaces’. *Topology* **26** (3) (1986) 311–335. With an appendix by M. Zisman. Cited on p. [5](#), [25](#)
- [BL87a] Brown, R. and Loday, J.-L. ‘Homotopical excision, and Hurewicz theorems for n -cubes of spaces’. *Proc. London Math. Soc. (3)* **54** (1) (1987) 176–192. Cited on p. [5](#)
- [BM99] Brown, R. and Mosa, G. H. ‘Double categories, 2-categories, thin structures and connections’. *Theory Appl. Categ.* **5** (1999) No. 7, 163–175 (electronic). Cited on p. [15](#)
- [BRS84] Brown, R. and Razak Salleh, A. ‘A van Kampen theorem for unions of nonconnected spaces’. *Arch. Math. (Basel)* **42** (1) (1984) 85–88. Cited on p. [9](#)
- [BS76a] Brown, R. and Spencer, C. B. ‘Double groupoids and crossed modules’. *Cahiers Topologie Géom. Différentielle* **17** (4) (1976) 343–362. Cited on p. [15](#), [16](#), [21](#)
- [BS76b] Brown, R. and Spencer, C. B. ‘ G -groupoids, crossed modules and the fundamental groupoid of a topological group’. *Nederl. Akad. Wetensch. Proc. Ser. A* **79**=*Indag. Math.* **38** (4) (1976) 296–302. Cited on p. [11](#), [14](#), [17](#)

- [CCHM] Cohen, C., Coquand, T., Huber, S. and Mörtberg, A. ‘Cubical Type Theory: a constructive interpretation of the univalence axiom’ *Leibniz International Proceedings in Informatics* 1–33 Schloss Dagstuhl Leibniz-Zentrum für Informatik. Cited on p. 16
- [Cro59] Crowell, R. H. ‘On the van Kampen theorem’. *Pacific J. Math.* **9** (1959) 43–50. Cited on p. 8, 9
- [CF63] Crowell, R.H. and Fox, R.H. *Introduction to Knot Theory*. Springer (1963). Cited on p. 7
- [Die89] Dieudonné, J. *A history of algebraic and differential topology. 1900–1960*. Birkhäuser Boston Inc., Boston-Basel, MA (1989). Cited on p. 7
- [Ehr65] Ehresmann, C. *Catégories et structures*. Dunod, Paris (1965). Cited on p. 11, 15, 16
- [Ein90] Einstein, A. ‘The Einstein-Wertheimer Correspondence on Geometric Proofs and Mathematical Puzzles’ *Math. Intell.* I, **12** (2)(1990) 35–43. Cited on p. 2
- [Ell93] Ellis, G. ‘Crossed squares and combinatorial homotopy’. *Math. Z.* **214** (1) (1993) 93–110. Cited on p. 6
- [ES87] Ellis, G. J. and Steiner, R. ‘Higher-dimensional crossed modules and the homotopy groups of $(n + 1)$ -ads’. *J. Pure Appl. Algebra* **46** (2-3) (1987) 117–136. Cited on p. 6, 25
- [Gro83] Grothendieck, A. *Pursuing stacks*, Manuscript, (1983) to appear in *Document Mathématique* édité par G. Maltsiniotis. Cited on p. 25
- [Gro] Grothendieck, A. *Correspondence with Ronald Brown*, edited M. Künzer to appear in *Document Mathématique*. available from <https://webusers.imj-prg.fr/~georges.maltsiniotis/ps.html> Cited on p. 10
- [Gro84] Grothendieck, A. *Esquisse d’un Programme* 54p. (1984). Cited on p. 23
- [Gou03] Goubault, E. ‘Some geometric perspectives in concurrency theory’. *Homology Homotopy Appl.* **5** (2) (2003) 95–136 (electronic). Algebraic topological methods in computer science (Stanford, CA, 2001). Cited on p. 16
- [Hig63] Higgins, P. J. ‘Algebras with a scheme of operators’. *Math. Nachr.* **27** (1963) 115–132. Cited on p. 15
- [Hig71] Higgins, P. J. *Notes on Categories and Groupoids*, Mathematical Studies, Volume 32. Van Nostrand Reinhold Co. (1971); Reprints in *Theory and Applications of Categories*, No. 7 (2005) pp 1-195. Cited on p. 9, 10
- [Hig05] Higgins, P. J. ‘Thin elements and commutative shells in cubical ω -categories’. *Theory Appl. Categ.* **14** (2005) No. 4, 60–74 (electronic). Cited on p. 14

- [Hu53] Hu, S.-T. ‘The homotopy addition theorem’. *Ann. of Math. (2)* **58** (1953) 108–122. Cited on p. 25
- [Jam] James, I.M. (editor) *History of Topology*, North-Holland (1994). Cited on p. 8
- [Kan55] Kan, D. M. ‘Abstract homotopy. I’. *Proc. Nat. Acad. Sci. U.S.A.* **41** (1955) 1092–1096. Cited on p. 16
- [Lod82] Loday, J.-L. ‘Spaces with finitely many nontrivial homotopy groups’. *J. Pure Appl. Algebra* **24** (2) (1982) 179–202. Cited on p. 25
- [MLW50] Mac Lane, S. and Whitehead, J. H. C. ‘On the 3-type of a complex’. *Proc. Nat. Acad. Sci. U. S. A.* **36** (1950) 41–48. Cited on p. 3
- [Mal09] Maltsiniotis, G. ‘La catégorie cubique avec connexions est une catégorie test stricte’. *Homology, Homotopy Appl.* **11** (2) (2009) 309–326. Cited on p. 16
- [Mas80] Massey, W. S. *Singular homology theory, Graduate Texts in Mathematics*, Volume 70. Springer-Verlag, New York (1980). Cited on p. 16
- [MV15] Munson, B.A. and Volić, I. *Cubical Homotopy Theory*, Cambridge Univ. Press (2015). Cited on p. 4
- [Pat12] Patchkoria, I. ‘Cubical approach to derived functors’ *Homology, Homotopy and Applications* **1** (2014) 133–158. Cited on p. 16
- [PR15] Paul, E. and Ramis, J.-P. ‘Dynamics on Wild Character Varieties’. *SIGMA* **11** (2015), 068, 21 pages, (electronic). Cited on p. 9
- [RZ16] Rivera, M. and Zeinalian, M. ‘Cubical rigidification, the cobar construction, and the based loop space’. *arXiv:1612.04801 [math.AT]* Cited on p. 16
- [Ste86] Steiner, R. ‘Resolutions of spaces by cubes of fibrations’. *J. London Math. Soc. (2)* **34** (1) (1986) 169–176. Cited on p. 25
- [Ton92] Tonks, A. ‘Cubical groups which are Kan’. *J. Pure Appl. Algebra* **81** (1) (1992) 83–87. Cited on p. 16
- [Vez14] Vezzani, A. ‘A motivic version of the theorem of Fontaine and Wintenberger’. *arXiv:1405.4548*. Cited on p. 16
- [Whi78] Whitehead, G. W. *Elements of homotopy theory, Graduate Texts in Mathematics*, Volume 61. Springer-Verlag, New York (1978). Cited on p. 25
- [Whi50] Whitehead, J. ‘A certain exact sequence’. *Ann. of Math. (2)* **52** (1950) 51–110. Cited on p. 24

- [Whi41] Whitehead, J. H. C. ‘On adding relations to homotopy groups’. *Ann. of Math. (2)* **42** (1941) 409–428. Cited on p. [17](#)
- [Whi49] Whitehead, J. H. C. ‘Combinatorial homotopy. II’. *Bull. Amer. Math. Soc.* **55** (1949) 453–496. Cited on p. [3](#), [5](#), [7](#), [17](#), [21](#), [22](#)
- [Zi72] Zisman, M. *Topologie Algébrique Élémentaire* Armand Collins (1972). Cited on p. [9](#)

Contact: www.groupoids.org.uk